

**EFFECT OF ECOLOGICAL CONDITIONS ON WOODY LITTER DECOMPOSITION  
PATTERN IN EIGHT WOODY SPECIES OF A DRY TROPICAL FOREST IN  
VINDHYAN HIGHLANDS**

**S.K. Pandey<sup>1</sup> and Hema Singh<sup>2</sup>**

<sup>1</sup>Govt. Art, Science and Commerce College, Jamgaon-R, Durg-491223 (C.G.), India  
Email: skpbhujnu2013@gmail.com

<sup>2</sup>Ecosystems Analysis Laboratory, Department of Botany, Banaras Hindu University, Varanasi  
221005 (India)

**ABSTRACT**

A study on litter decomposition of eight woody species, viz. *Shorea robusta*, *Buchanania lanzan*, *Diospyros melanoxylon*, *Lagerstroemia parviflora*, *Lannea coromandelica*, *Terminalia tomentosa*, *Holarrhena antidysenterica* and *Lantana camara* was done. Litter decomposition was monthly monitored using litter bag technique over a 12-month period in a dry tropical deciduous forest of Vindhyan highland, India. Wood litter weight loss ranged from 14.08% in *T. tomentosa* at Kotwa site to 25.25% in *T. tomentosa* at Hathinala site and month wise wood litter weight loss ranged from 0.60% in January for *L. parviflora* to 47.53% in August for *L. coromandelica*. The chemical composition of litter affected decomposition pattern. Nitrogen and phosphorus mineralization rates and decay pattern varied significantly from species to species. *T. tomentosa* having higher nitrogen content and lower C/N ratio exhibited faster weight loss than other species such as *S. robusta*, *D. melanoxylon*. The initial nitrogen, phosphorus contents of litter showed significant positive relation with weight loss in monthly retrieved litter. C/N ratio showed significant negative relationship with decay constant. Soil temperature and soil moisture had positive relation with weight loss. Our results suggest that dry tropical deciduous forest species are characterized by faster decomposition, indicating a faster rate of organic matter turnover and rapid nutrient cycling due to intra-annual changes in the environmental conditions. This could influence the soil nutrient content in system.

**Keywords** Litter decomposition, environmental, weight loss, woody species, C/N ratio, decay constant, dry tropical deciduous.

**INTRODUCTION**

Litter decomposition is an important process that connects many aboveground and belowground processes. The decomposition of dead leaves and roots is one of the major pathways by which  
© S. K.Pandey et al. J. Sci. Res. Adv. Vol. 3, No. 1, 2016, 235-251| 235

carbon fixed during photosynthesis is returned to the atmosphere (Coûteaux et al., 1995). The process also contributes to soil. Decomposing plant litter and soil organic matter play an important role in the terrestrial global carbon cycle (Schimel, 1995). Decomposition of leaf litter, by which organic matter and nutrients are returned to the forest soils, has received considerable attention for sustainable soil fertility (Swift et al., 1979; Hobbie 1992; Moretto et al., 2001; Xuluc-Tolosa et al., 2003; Moore et al., 2006; Sayer 2006). The rate of decomposition depends on environmental controls such as climate, litter quality and soil characteristics. Studies have indicated association of the decomposition with the carbon and nitrogen contents of the litter (Fog, 1988; Kemp et al., 2003; Meentemeyer, 1978; Swift et al., 1979). Complexes of bacteria, fungi, and soil organisms have an important role in decomposing leaf litter. The micro-fauna, which can move freely through the litter bag net, have been shown to cause an increase in weight loss (Berg et al., 1980; Berg and Staaf, 1981). Plant detritus and soil organic carbon are the largest carbon pools in terrestrial biosphere (Moore and Braswell, 1994), and therefore, understanding litter decomposition processes and the factors controlling litter decomposition is important for studying nutrient cycling, developing carbon budgets as well as assessing implications of global climate change. Litter decomposition rates are a function of litter quality, biota and microclimate (Seastedt et al., 1983), as well as edaphic properties (Heneghan et al., 1998). The successional changes of soil organisms have been demonstrated during the decomposition process of various litters using the litter bag method (Berg and Soderstrom, 1979). The roles of soil organisms in nutrient cycle may change during the decomposition processes of litter (Hasegawa and Takeda, 1995; Warren and Zou, 2002). In decreasing order of importance, the key factors regulating decomposition are commonly assumed to be climate, litter quality (e.g. N content, C/N ratio) and decomposer communities (e.g. bacteria, fungi, and soil macro- and micro fauna) (Meentemeyer 1984). Many studies have concluded that the combination of climate (e.g. mean annual temperature, mean annual precipitation etc) and litter quality (N content, C: N ratio) are the primary factors controlling litter decomposition (Coûteaux et al. 1995; Aerts, 1997; Moorhead et al. 1999; Gholz et al. 2000; Silver and Miya 2001). Across a wide variety of litter type C/N ratio seems to be top interpreter of decay rate (Perez-Harguindeguy et al. 2000). The significance of litter decomposition to nutrient cycling and ecosystem role has been known, and a several studies has been undertaken (Gillon et al. 1999; Moore et al. 2006), various in response to provincial or species-specific concerns. Attempts to calculate the effects of climate change have newly driven studies of litter from broad ranges of plant types (Perez-Harguindeguy et al. 2000). The effect of litter quality is observable from the different decay rates of different tissue types, but identifying the particular litter characteristics that are every time and directly associated to decomposability has verified unpredictably complicated. Different species have dissimilar nutrient discharge patterns, which are linked to quality, time of year, and ecological factors (Abiven et al., 2005; Arunachalam et al., 2003). The Vindhyan highlands that are covered by dry tropical deciduous forest have high woody species diversity (Jha and Singh 1990; Sagar and Singh 2004). However, there is too small information on the comparative rates of decay of leaf and wood litter of these species and litter chemical parameters helpful for predicting their decompose rates in dry tropical deciduous forest. Thus, in this study we have quantified the monthly woody litter decomposition rates and nutrient

dynamics of the dominant dry tropical deciduous forest woody species: *S. robusta*, *B. lanzan*, *D. melanoxyton*, *L. parviflora*, *L. coromandelica*, *T. tomentosa*, *H. antidysenterica* and *L. camara* of Vindhyan highlands, Uttar Pradesh, India, to know their role in the organic matter and nutrient returns that has a bearing on the total productivity of the ecosystem.

## MATERIALS AND METHODS

### *Study site*

The study was conducted on four sites, viz. Kotwa (25°00'17'' N and 82°37'38'' E, 196 m.a.s.l.), Ranitalli (24°18'11'' N and 83°04'22'' E, 287 m.a.s.l.), Neruiadamer (24°18'33'' N and 83°23'05'' E, 323 m.a.s.l.), Bokarakhari (24°24'13'' N and 83°12'01'' E, 245 m.a.s.l.) and Hathinala (24°18'07'' N and 83°05'57'' E, 291 m.a.s.l.) of the Vindhyan dry tropical region, Sonbhadra District and Kotwa in Mirzapur district, Uttar Pradesh (Fig. 1). The area experiences a tropical monsoon climate. The sites are located between two meteorological stations, Obra and Renukut. Ranitalli site is nearest to Obra and Hathinala site is nearest to Renukut. Mean annual rainfall is 926 mm at Obra and 1146 mm at Renukut (Pandey et al., 1992). The soils are residual ultisols, sandy loam in texture, and reddish to dark grey in colour and are extremely poor in nutrients (Singh et al., 1989). Physio-chemical property of five experimental sites (Table 1). The mean monthly site wise soil moisture content (%) was shown in Fig. 2.

### *Litter decomposition*

Woody litter decomposition experiment was carried out during a 12-month period. The collected litter was spread separately in thin layer for drying at room temperature (25 – 30°C). In order to study the effect of environmental conditions on the rate of decomposition, litter bags were placed at the forest floor of each month and recovered at the end of each month. Thus, results in twelve times within the year. Litter decomposition studies were carried out by using the litter bag (15 × 15 cm) technique (Gilbert and Bockock, 1960). Nylon net litter bags, each containing 10 gm of air dried leaf litter, were placed on the forest floor at each site to study the percentage weight loss. The mesh size of the bags was 1 mm which easily allows the microbial activity of soil microflora and fauna. 288 litter bags were used in which 24 litter bags (i.e. three replicate for each species) placed on the soil surface in each month; each site and three replicate litter bags were sampled at each month for each species at each site for wood litter weight loss. The bags were sealed with stainless steel staples and labeled with aluminum tags. Each litter bag was attached to the ground by nylon wire to prevent movement and ensure good contact between the bags and the surface layer on experimental sites. Each litter bag collected was put on individual bag and transported to the laboratory. Before the measurement of litter weights, litter was cleared of any adhering plant and soil fragments. Weights of wood litter were recorded before and after oven drying to constant weight at 80 °C. After the measurement of weights, wood litter samples were taken for chemical analysis. Carbon and nitrogen estimated by CHN analyser and phosphorus by phosphomolybdic blue colorimetric method (Anderson, 1989). Litter weight losses for eight

woody plant species (Table 2), during twelve months of the study period were calculated using the following formula:

$$[\text{Weight loss (\%)} = (\text{Initial wt.} - \text{Final wt.}) / (\text{Initial wt.}) \times 100]$$

### **Decay constant (k)**

The monthly decay constant (k/mo) for wood litter weight loss was calculated by following the negative exponential decay model (Olson 1963, Ezcurra, E., and J. Becerra. 1987):  $k = -\ln(X/X_0)/t$ , where,  $X_0$  is the initial dry weight,  $X$  is the dry weight remaining at the end of the investigation, and  $t$  is the time period (month or year). The time required for 50% ( $t_{50}$ ) and 95% ( $t_{95}$ ) weight loss was calculated as  $t_{50} = 0.693/k$  and  $t_{95} = 3/k$ .

## RESULTS

The coefficient of variation for wood litter weight loss was maximum for *S. robusta* and *D. melanoxylon* (90%) followed by *H. antidysenterica* (89%), *L. coromandelica* (88%), *L. parviflora* (87%), *B. lanzan* (86%), *L. camara* (85%) and minimum for *T. tomentosa* (82%), which had maximum wood weight loss. Wood litter weight loss of all species also positively related to soil temperature and soil moisture as in Fig. 3. Analysis of variance indicated that wood litter weight loss of all species had significant site effect. *S. robusta* ( $F_{4, 120} = 10.84, P = 0.000$ ), *B. lanzan* ( $F_{4, 120} = 64.46, P = 0.000$ ), *D. melanoxylon* ( $F_{4, 120} = 13.13, P = 0.000$ ), *L. parviflora* ( $F_{4, 120} = 9.34, P = 0.000$ ), *L. coromandelica* ( $F_{4, 120} = 8.72, P = 0.000$ ), *T. tomentosa* ( $F_{4, 120} = 81.72, P = 0.000$ ), *H. antidysenterica* ( $F_{4, 120} = 25.54, P = 0.000$ ) and *L. camara* ( $F_{4, 120} = 18.35, P = 0.000$ ). ANOVA also indicated that wood litter weight loss of all species had significant site  $\times$  month effect. *S. robusta* ( $F_{44, 120} = 4.47, P = 0.000$ ), *B. lanzan* ( $F_{44, 120} = 6.40, P = 0.000$ ), *D. melanoxylon* ( $F_{44, 120} = 4.58, P = 0.000$ ), *L. parviflora* ( $F_{44, 120} = 1.80, P = 0.000$ ), *L. coromandelica* ( $F_{44, 120} = 3.51, P = 0.000$ ), *T. tomentosa* ( $F_{44, 120} = 5.00, P = 0.000$ ), *H. antidysenterica* ( $F_{44, 120} = 6.64, P = 0.000$ ), and *L. camara* ( $F_{44, 120} = 3.28, P = 0.000$ ). Mean carbon content of decomposed woody litter varied from  $37.07 \pm 0.01\%$  in August to  $38.07 \pm 0.01\%$  in December and January, mean nitrogen content varied from  $0.75 \pm 0.01\%$  in January to  $0.80 \pm 0.003\%$  in July and August, mean phosphorus content varied from  $0.085 \pm 0.001\%$  in January to  $0.09 \pm 0.00\%$  in July and mean C/N ratio varied from 47.01 in August to 50.76 in January in Fig. 4. Site wise variation in wood litter weight loss at five sites varied from 14.08% in *T. tomentosa* at Kotwa site to 25.25% in *T. tomentosa* at Hathinala site. Month wise wood litter weight loss ranged from 0.60% in January for *L. parviflora* to 47.53% in August for *L. coromandelica* in Fig. 5. C/N ratio of monthly retrieved woody litter weight loss minimum in July and August; maximum in January and December.

### **Decay constant (k)**

The coefficient of variation for  $k$  was 89% for wood litter weight loss. ANOVA also indicated a significant site effect ( $F_{4, 960} = 7220, P = 0.000$ ), species ( $F_{7, 960} = 1462, P = 0.000$ ) and month wise effect ( $F_{11, 960} = 229900, P = 0.000$ ). Monthly decay constant of wood litter varied from 0.2259 at Kotwa site to 0.3041 at Hathinala site (1.3-fold), species wise decay constant varied

from 0.2332 for *H. antidysenterica* to 0.2834 for *S. robusta* (1.2-fold) and month wise decay constant varied from 0.0116 in January to 0.5998 in August (59-fold) (Table 3 and 4).

## DISCUSSION

In this study we find that wood litter weight loss for different tree species were positively related with soil temperature and soil moisture content (Fig. 1). During decomposition, decay of the organic nitrogen into an inorganic form mainly depends on the C/N ratio of the material. The initial weight loss is promoted by high nitrogen and low C/N ratio (Edmonds, 1990, Jamaludheen and Kumar, 1999). Taylor et al., (1989), found that C/N ratio to be a better interpreter of weight loss than the lignin/N ratio in a microcosm decomposition study of leaf litter. C/N ratio of woody litter was more than 25 for all species. This validates the high quality of their litter as Myers et al., (1994), reported that substrate with C/N ratio is less than 25 is of high quality and releases N at faster rate than low quality residues (C/N > 25). Mean carbon content and C/N ratio of decomposed litter decreased while nitrogen and phosphorus increased, in our results. Carbon concentration was negatively related with nitrogen and phosphorus concentration in decomposed litter and showed positive relation with C/N ratio. N and P concentrations in the decomposing wood litter of different plant species were highly changeable and followed a similar pattern during the decay process. Along with this, N and P concentration in decomposing litter increased constantly until the end of the study, the degree of increase, however, differed among the species (Fig. 4), as Singh et al., 1999 also reported. Our result showed that increase of the weight loss from the mid of the experiment in *T. tomentosa*, *L. coromendelica*, due to high N content (Fig. 5). However, the slow rate of initial decay was followed by a rapid phase of decomposition, may be because of the quicker breakdown of litter by the decomposing population that initially took time to colonize the litter (Songwe et al., 1995) and slower decay at the later stage could, however, be due to the accumulation of more unbreakable materials as shown in our results. The changeable rate of decomposition seemed to be because of not only to the nutrient content of the leaves but also to their morphological characteristics (Songwe et a., 1995). The decrease of carbon content with time must be related to the carbohydrates and phenolic compounds degradation at the beginning of the process (Schlesinger 1985), which is also related to mineralization in humid periods (Santa Regina, San Miguel, and Gallardo, 1986). When the litter is exposed to decomposition microorganisms, the easily decomposable components are attacked earliest. The soluble and polymer carbohydrates (cellulose and hemicelluloses) disappearing as faster than the other (Hirobe et al., 2004a, b). Many studies have also supported that litter with high N concentrations and a low C/N ratio is decomposed more rapidly than those with low N concentrations and a high C/N ratio (Finzi et al., 1998; Berg and Laskowski, 2006; Kamei et al., 2009). In this study lower mass loss during winter, summer season and faster in rainy season may be due to cold and dry conditions in winter, and warm and humid condition in rainy season , respectively; also reported by Sangha et al., 2006, Arunachalam et al., 1998, this also in dry season Mathani et al., 1996). Litter mass loss showed positive correlation with rainfall and soil moisture (Dasseler et al., 2000; Okeke et al., 1992; Austin et al., 2000). Soil nutrient content could not be related to litter mass loss, which is in agreement with Prescott (1995), Aerts and De

Caluwe (1997), who has indicated that N availability did not alter litter decomposition rates in forests. These suggest that a small amount of the N release from litter was mineralized by soil microbes (Prescott, 1995). The increase in N concentration in all species is a worldwide study in decomposing litter (Mellilo et al., 1982; Moro and Domingo, 2000; Songwe et al., 1995). Woody litter decomposed faster, which may be due to decomposition of wood by termite activity. That was observed during our study period that termites attacked woody litter robustly. Termites accelerate the litter decomposition of tropical forest which is an important faunal component (Peterson et al., 1982; Yamashita et al., 1998; Sandhu et al., 1990 and Swift et al., 1979). Phosphorus mineralization during decomposition was quite similar to that of N release; the rate of release was slightly slower than that of nitrogen (Berg and Staff 1980). Subsequent increase in N and P concentrations could be the result of microbial immobilization (Maithani et al., 1996), nutrient inputs from through fall, and atmospheric precipitation (Bocock, 1963) and or atmospheric N fixation (Wood, 1974) as also observed in our leaf and wood litter decomposition study. Annual decay constant varied from 1.16 in *S. robusta* to 1.53 in *T. tomentosa*, which was maximum (Table 3). While Roy and Singh reported (1994)  $k = 1.93-2.26$  for dry tropical deciduous forest litter. Therefore, our range is lower than this but supported the range  $k = 0.64-1.30$ , reported by K.P. Singh et al., 1999. Moisture availability in the form of rainfall is naturally a key climate state factor that directly or indirectly controls the construction and function of tropical forest ecosystems since seasonal variation in temperature tends to be small (Richards and Caldwell 1987; Clark et al., 2003; Feeley et al., 2007). Many studies also supported that litter with high nitrogen content decomposed more rapidly (Kamei et al., 2009; Tripathi and Singh, 1992a). The decomposition activity is driven by the decomposer (Tian et al., 1997, Tian and Tateishi, 2002). The activity is also regulated by the nutrient availability and carbon energy resources.

## CONCLUSION

On the whole, it can be concluded that the nutrient in decomposing wood litter are recycled rapidly within the forest soil surface. Litter weight loss was relatively high and differed significantly among the species and on soil temperature and soil moisture condition. Both initial nitrogen and phosphorus concentration had positive influence on decomposition while litter decomposition of eight species was negative relationship with C/N ratio. Therefore, it is obvious that initial nutrients concentrations along with soil moisture and temperature take part in determining the litter decomposition in the dry tropical deciduous forest and thus organic matter turnover, nutrient cycling in the forest.

## ACKNOWLEDGEMENT

We thank the Ministry of Environment and Forests, Government of India for providing financial assistance.

**Table 1.** Physio-chemical properties of the five experimental sites located in Vindhyan highlands (Mean $\pm$ SE).

Parameters	Kotwa	Ranitalli	Neruiadamer	Bokarakhari	Hathinala
Sand (%)	70.3±0.89	72.0±1.36	67.7±0.93	65.4±1.58	63.0±0.93
Silt (%)	26.9±1.20	26.0±1.91	28.1±0.95	27.9±1.78	26.3±0.98
Clay (%)	02.1±0.50	2.0±0.79	4.2±0.63	6.7±1.21	10.1±1.73
pH	07.0±0.10	7.1±0.17	6.9±0.23	6.4±0.13	6.5±0.18
Total-C (%)	1.20±0.10	1.3±0.15	1.2±0.27	1.4±0.29	1.5±0.38
Total-N (%)	0.11±0.00	0.14±0.03	0.13±0.01	0.11±0.01	0.13±0.02
Total-P (%)	0.02±0.00	0.02±0.01	0.03±0.02	0.02±0.01	0.05±0.02
C/N ratio	10.91	9.29	9.23	12.73	11.54

**Table 2.** List of the species for decomposition study of five experimental sites located in Vindhyan highlands.

Name of the species	Common name	Family	Growth form
<i>Shorea robusta</i> Gaertn. f.	Sal	Dipterocarpaceae	Tree
<i>Buchanania lanzan</i> Spreng.	Piyar	Anacardiaceae	Tree
<i>Diospyros melanoxylon</i> Roxb.	Tendu	Ebenaceae	Tree
<i>Lagerstroemia parviflora</i> Roxb.	Siddh	Lythraceae	Tree
<i>Lannea coromandelica</i> L.	Jigan	Anacardiaceae	Tree
<i>Terminalia tomentosa</i> (Roxb.) Wight & Arn.	Asna	Combrataceae	Tree
<i>Holarrhena antidysenterica</i> Wall.	Koraya	Apocynaceae	Tree
<i>Lantana camara</i> L.	Lantana	Verbenaceae	Shrub

Nomenclature based on 'Flora of Madhya Pradesh' 1997 (eds. Mudgal V, Khanna KK, Hajra PK) Botanical Survey of India, Calcutta, India.

**Table 3.** Mean decay constant (monthly) of different woody litter species on different sites.

Site*	k
Kotwa	0.2259 <sup>a</sup>
Ranitalli	0.2656 <sup>b</sup>
Neruiadamar	0.2753 <sup>c</sup>
Bokarakhari	0.2840 <sup>d</sup>
Hathinala	0.3041 <sup>e</sup>
Species**	
<i>S. robusta</i>	0.2834 <sup>a</sup>
<i>B. lanzan</i>	0.2834 <sup>a</sup>
<i>D. melanoxylon</i>	0.2752 <sup>b</sup>
<i>L. parviflora</i>	0.2782 <sup>c</sup>
<i>L. coromandelica</i>	0.2789 <sup>c</sup>
<i>T. tomentosa</i>	0.2692 <sup>d</sup>
<i>H. antidysenterica</i>	0.2332 <sup>e</sup>
<i>L. camara</i>	0.2663 <sup>f</sup>

\*Values are across species and months


\*\*Values are across sites and months

Values suffixed with different letters for sites and species are significantly different from each other at  $P < 0.05$ .

**Table 4.** Monthly decay constants of woody litter during a one year cycle in Vindhyan highlands.

Month	Decay constant k (/mo)
January	0.0116
February	0.0299
March	0.0524
April	0.0591
May	0.0896
June	0.4468
July	0.5710
August	0.5998
September	0.5917
October	0.5267
November	0.2355
December	0.0375





**Figure 1.** Map showing the study sites.







Figure 2. Monthly soil moisture content (SMC) at five experimental sites.



**Figure 3.** Monthly woody weight loss (%), soil temperature



**Figure 4** **A.** Monthly woody litter weight loss (%) and litter carbon content (%). **B.** Monthly woody litter weight loss (%) and litter phosphorus content. **C.** Monthly woody litter weight loss (%) and litter nitrogen content. **D.** Monthly pattern woody litter weight loss (%) and C/N ratio.



**Figure 5.** Decomposition pattern of woody litter as exhibited by weight loss of eight species at five sites: A, *S. robusta*; B, *B. lanzan*; C, *D. melanoxylon*; D, *L. parviflora*; E, *L. coromandelica*; F, *T. tomentosa*; G, *H. antidysenterica*; H, *L. camara*.

REFERENCES

1. Abiven, S., Recous S., Reyes V. and Oliver R. 2005. Mineralisation of C and N from root, stem and leaf residues in soil and role of their biochemical quality. *Biol Fert Soils* **42**: 119–128.
2. Aerts, R. and De Caluwe H. 1997. Nutritional and plant-mediated controls on leaf litter decomposition of *Carex* species. *Ecology* **78**: 244–260.
3. Aerts, R. 1997. Climate, leaf litter chemistry and leaf litter decomposition in terrestrial ecosystems: a triangular relationship. *Oikos* **79**: 439–449.
4. Anderson, J.M. and Ingram J.S.I. 1989. In *Tropical Soil Biology and Fertility: A Handbook of Methods of Analysis*, CAB International, Wallingford, UK, 1st edn.
5. Arunachalam, A., Maithani K., Pandey H.N. and Tripathi R.S. 1998. Leaf litter decomposition and nutrient mineralization patterns in regrowing stands of a humid subtropical forest after tree cutting. *For Ecol Manage* **109**: 151–161.
6. Arunachalam, K., Singh N.D. and Arunachalam A. 2003. Decomposition of leguminous crop residues in jhum cultivation system in northeast India. *J Plant Nutr Soil Sci* **166**: 731–736.
7. Berg, B. and Soderstrom B. 1979. Fungal biomass and nitrogen in decomposing Scot pine needle litter. *Soil Biology and Biochemistry* **11**: 339–341.
8. Berg, B. and Staaf H. 1981. Leaching, accumulation and release of nitrogen in decomposing forest litter. *Ecological Bulletins* **33**: 163–178.
9. Berg, B. and Staff H. 1980. Decomposition rate and chemical changes in Scot pine needle litter. 2. Influence of chemical composition. *Ecological Bulletins* **32**: 363–372.
10. Berg, B., Lohm U., Lundkvist H. and Wiren A. 1980. Influence of soil animals on decomposition of Scots pine needle litter. *Ecological Bulletins* **32**: 401–409.
11. Bockock, K.L., Gilbert O.J., Capstick C.K., Turner D.C., Ward J.S. and Woodman M.J. 1960. Changes in leaf litter when placed on the surface of soil with contrasting humus types. *Journal of Soil Science* **11**: 1-9.
12. Bradstreet, R.B. 1965. The Kjeldahl Method for Organic Nitrogen, Academic Press, New York, p. 239.
13. Clark, D.A., Piper S.C., Keeling C.D. and Clark D.B. 2003. Tropical rain forest tree growth and atmospheric carbon dynamics linked to interannual temperature variation during 1984–2000. *Proc Natl Acad Sci* **100**: 5852–5857.
14. Coûteaux, M.M., Bottner P. and Berg B. 1995. Litter decomposition, climate and litter quality. *Trends Ecological Evolution* **10**: 63–66.
15. Edmonds, R.L. 1990. Litter decomposition and nutrient release in Douglas fir, red alder, western hemlock and Pacific silver fir ecosystems in western Washington. *Canadian Journal of Forest Research* **10**: 327–337.
16. Ezcurra, E. and Becerra J. 1987. Experimental decomposition of litter from the Tamaulipan Cloud Forest: A comparison of four simple models. *Biotropica* **19**: 290–296.
17. Feeley K.J., Joseph Wright S., Nur Supardi M.N., Kassim A.R., Davies S.J. 2007. Decelerating growth in tropical forest trees. *Ecol Lett* **10**: 461–469.

18. Fog, K. 1988. The effect of added nitrogen on the rate of decomposition of organic matter. *Biological Review* **63**: 433–462.
19. Gholz, H.L., Wedin D.A., Smitherman S.M., Harmon M.E. and Parton W.J. 2000. Long-term dynamics of pine and hardwood litter in contrasting environments: toward a global model of decomposition. *Global Change Biology* **6**: 751–765.
20. Gillon, D., Joffre R. and Ibrahima A. 1999. Can litter decomposability be predicted by near infrared reflectance spectroscopy? *Ecology* **80**: 175–186.
21. Hasegawa, M. and Takeda H. 1995. Changes in feeding attributes of four collembolan populations during the decomposition process of pine needles. *Pedobiologia* **39**: 155–169.
22. Heneghan, L., Coleman, D.C., Zou, X., Jr. Crossley D.A. and Haines B.L. 1998. Soil microarthropod community structure and litter decomposition dynamics: a study of tropical and temperate sites. *Applied Soil Ecology* **9**: 33–38.
23. Hirobe, M., Sabang J., Bhatta B.K. and Takeda H. 2004a. Leaf-litter decomposition of 15 tree species in a lowland tropical rain forest in Sarawak: decomposition rates and initial litter chemistry. *J For Res* **9**: 341–346.
24. Hirobe, M., Sabang J., Bhatta B.K. and Takeda H. 2004b. Leaf-litter decomposition of 15 tree species in a lowland tropical rain forest in Sarawak: dynamics of carbon, nutrients, and organic constituents. *J For Res* **9**: 347–354.
25. Hobbie, S.E. 1992. Effects of plant-species on nutrient cycling. *Trends Ecol Evol* **7**: 336–339.
26. Jamaludheen, V. and Kumar B.M. 1999. Litter of multipurpose trees in Kerala, India: Variations in the amount, quality, decay rates and release of nutrients. *Forest Ecology and Management* **115**: 1–11.
27. Jha, C.S. and Singh J.S. 1990. Composition and dynamics of dry tropical forest in relation to soil texture. *Journal of Vegetation Science* **1**: 609–614.
28. Kemp, P.R., Reynolds, J.F., Virginia R. and Whitford W.J. 2003. Decomposition of leaf and root litter of Chihuahuan desert shrubs: effect of three years of summer drought. *Journal of Arid Environment* **53**: 21–39.
29. Maithani, K., Tripathi R.S., Arunachalam A. and Pandey H.N. 1996. Seasonal dynamics of microbial biomass C, N and P during regrowth of a disturbed subtropical humid forest in northeast India. *Appl Soil Ecol* **4**: 31–37.
30. Meentemeyer, V. 1978. Macroclimate and lignin control of litter decomposition rates. *Ecology* **59**: 465–472.
31. Meentemeyer, V. 1984. The geography of organic decomposition rates. *Annals of the Association of American Geographers* **74**: 551–560.
32. Melillo, J.M, Aber J.B. and Muratore J.F. 1982. Nitrogen and lignin control of hardwood leaf litter decomposition dynamics. *Ecology* **63**: 621–626.
33. Moore T.R., Trofymow J.A., Prescott C.E., Fyles J. and Titus B.D. 2006. Patterns of carbon, nitrogen and phosphorus dynamics in decomposing foliar litter in Canadian forests. *Ecosystems* **9**: 46–62.

34. Moorhead, D.L., Currie W.S., Rastetter E.B., Parton W.J. and Harmon M.E. 1999. Climate and litter quality controls on decomposition: an analysis of modeling approaches. *Global Biogeochemical Cycles* **13**: 575–589.
35. Moretto, A.S., Distel R.A. and Didone N.G. 2001. Decomposition and nutrient dynamic of leaf litter and roots from palatable and unpalatable grasses in a semi-arid grassland. *Applied Soil Ecology* **18**: 31-37.
36. Moro, M.J. and Domingo F. 2000. Litter decomposition in four woody species in a Mediterranean climate: weight loss, N and P dynamics. *Ann. Bot.* **86**: 1065–1071.
37. Myers, R.J.K., Palm C.A., Cuevas E., Gunatilleke I.U.N. and Brossard M. 1994. The synchronization of nutrient mineralization and plantnutrient demand. In: Woomer PL, Swift MJ (eds) *The biological management of tropical soils fertility*. A Wiley-Sayce Publication, Chichester, pp 81–116.
38. Negi, J.D.S., Chauhan P.S. and Negi M. 2003. *Indian Forester* **129**: 757–769.
39. Okeke, A. I. and Omaliko C.P.E. 1992. Leaf litter decomposition and carbondioxide evolution of some agroforestry fallow species in southern Nigeria. *Forest Ecology and Management* **50**: 103–116.
40. Olson, J.S. 1963. Energy storage and the balance of producers and decomposers in ecological systems. *Ecology* **44**: 322–331.
41. Pandey, C.B. and Singh J.S. 1992. Rainfall and grazing effects on net primary productivity in a tropical savanna, India. *Ecology* **73**: 2007–2021.
42. Perez-Harguindeguy, N., Diaz S., Cornelissen J.H.C., Vendramini F., Cabido M. and Castellanos A. 2000. Chemistry and toughness predict leaf litter decomposition rates over a wide spectrum of functional types and taxa in central Argentina. *Plant Soil* **218**: 21–30.
43. Peterson, H. and Luxton M. 1982. A comparative analysis of soil fauna populations and their role in decomposition processes. *Oikos* **39**: 288–388.
44. Prescott, C.E. 1995. Does nitrogen availability control rates of litter decomposition in forest. *Plant and Soil* **168–169**: 83–88.
45. Richards, J.H. and Caldwell M.M. 1987. Hydraulic lift: substantial nocturnal water transport between soil layers by *Artemesia tridentata* roots. *Oecologia* **73**: 486–489.
46. Roy, S. and Singh J.S., 1994. Consequences of habitat heterogeneity for availability of nutrients in a dry tropical forest. *J. Ecol.* **82**: 503-509.
47. Sagar, R. and Singh J.S. 2004. Local plant species depletion in a tropical dry deciduous forest of northern India. *Environmental Conservation* **31**: 55-62.
48. Sandhu, J., Sinha M. and Ambasht R.S. 1990. Nitrogen release from decomposing litter of *Leuceana leucocephala* in the dry tropics. *Soil Biology and Biochemistry* **22**: 859–863.
49. Sangha, K.K., Jalota R.K. and Midmore D.J. 2006. Litter production, decomposition and nutrient release in cleared and uncleared pasture systems of central Queensland, Australia. *J Trop Ecol* **22**: 177–189.
50. Santa Regina, I., San Miguel C. and Gallardo F.J. 1986. Evolución y velocidad de descomposición de la hojarasca en tres bosques de la Sierra de Bejar (Salamanca). *Anuario CEBAS CSIC Salamanca* **11**: 217–231.

51. Sayer, E.J. 2006. Using experimental manipulation to assess the roles of leaf litter in the functioning of forest ecosystems. *Biol Rev* **81**: 1–31.
52. Schimel, D.S. 1995. Terrestrial ecosystems and the carbon cycle. *Global Change Biology* **1**: 77–91.
53. Schlesinger, W.H. 1985. Decomposition of chaparral shrub foliage. *Ecology* **66** (4): 1353–1359.
54. Seastedt, T.R., Crossley D.A., Jr. Meentemeyer V. and Waide J.B. 1983. A two-year study of leaf litter decomposition as related to macroclimatic factors and microarthropod abundance in the southern Appalachians. *Holarctic Ecology* **6**: 11–16.
55. Silver, W.L. and Miya R.K. 2001. Global patterns in root decomposition: comparisons of climate and litter quality. *Oecologia* **129**: 407–419.
56. Singh, J.S., Raghubanshi A.S., Singh R.S. and Srivastava S.C. 1989. Microbial biomass acts as a source of plant nutrient in dry tropical forest and savanna. *Nature* **338**: 499–500.
57. Singh, J.S., Raghubanshi A.S., Singh R.S. and Srivastava S.C. 1989. Microbial biomass acts as a source of plant nutrient in dry tropical forest and savanna. *Nature* **338**: 499–500.
58. Singh, K.P., Singh P.K. and Tripathi S.K. 1999. Litterfall, litter decomposition and nutrient release patterns in four native tree species raised on coal mine spoil at Singrauli, India. *Biol Fertil. Soils* **29**: 371–378.
59. Songwe, N.C., Okali D.U.U. and Fasehun F.E. 1995. Litter decomposition and nutrient release in a tropical rainforest, Southern Bakundu Forest Reserve, Cameroon. *J. Trop. Ecol.* **11**: 333–350.
60. Swift, M.J., Heal O.W. and Anderson J.M. 1979. Decomposition in terrestrial ecosystems. *Studies in Ecology, vol. 5. University of California Press Berkeley, CA.*
61. Swift, M.J., Heal O.W. and Anderson J.M. 1979. In *Decomposition in Terrestrial Ecosystems, Blackwell Scientific Publications, UK.*
62. Taylor, B., Parkinson B., Parsons W.F.J. 1989. Nitrogen and lignin control of hardwood leaf decomposition dynamics. *Ecology* **63**: 621–626.
63. Tian, X., Takeda H. and Ando T. 1997. Application of a rapid thin section method for observations on the decomposing litter in the more humus form in a subalpine coniferous forest. *Ecol Res* **12**: 289–300.
64. Tian, X-J. and Tateishi T. 2002. Relative roles of microorganisms and soil animals on needle litter decomposition in a subalpine coniferous forest. *Acta Phytoecol Sin* **26**: 257–263.
65. Tripathi, S.K. and Singh K.P. 1992a. Abiotic and litter quality control during decomposition of different plant parts in a dry tropical bamboo savanna in India. *Pedobiologia* **36**: 241–256.
66. Warren, M.W. and Zou X. 2002. Soil macrofauna and litter nutrients in three tropical tree plantations on a disturbed site in Puerto Rico. *Forest Ecology and Management* **170**: 161–171.
67. Xuluc-Tolosa, F.J., Vester H.F.M., Ramí ´rez-Marcial N., Castella-nos-Albores J. and Lawrence D. 2003. Leaf litter decomposition of tree species in three successional phases



- of tropical dry secondary forest in Campeche, Mexico. *Forest Ecology and Management* **174**: 401–412.
68. Yamashita, T. and Takeda H. 1998. Decomposition and nutrient dynamics of leaf litter in litter bags of two mesh sizes set in two Dipterocarpus forest sites in peninsular Malaysia. *Pedobiologia* **42**: 11–21.